

Publications

Valentonyte, R. *et al.* Sarcoidosis is associated with a truncating splice site mutation in BTNL2. *Nature genetics* **37**, 357–64 (2005).

Augustin, R. *et al.* Dickkopf related genes are components of the positional value gradient in Hydra. *Dev Biol* **296**, 62–70 (2006).

Franke, A. *et al.* No association between the functional CARD4 insertion/deletion polymorphism and inflammatory bowel diseases in the German population. *Gut* **55**, 1679–1680 (2006).

Franke, A. et al. GENOMIZER: an integrated analysis system for genome-wide association data. Hum Mutat 27, 583–588 (2006).

Schafmayer, C. et al. Investigation of the Lith1 candidate genes ABCB11 and LXRA in human gallstone disease. Hepatology 44, 650–657 (2006).

Buch, S. et al. A genome-wide association scan identifies the hepatic cholesterol transporter ABCG8 as a susceptibility factor for human gallstone disease. Nature genetics 39, 995–9 (2007).

Franke, A. et al. Systematic association mapping identifies NELL1 as a novel IBD disease gene. PLoS ONE 2, e691 (2007).

Franke, A. et al. No association between the TUCAN (CARD8) Cys10Stop mutation and inflammatory bowel disease in a large retrospective German and a clinically well-characterized Norwegian sample. Gastroenterology 132, 2080–1 (2007).

Hampe, J. et al. A genome-wide association scan of nonsynonymous SNPs identifies a susceptibility variant for Crohn disease in ATG16L1. Nature genetics 39, 207–11 (2007).

Prescott, N. J. et al. A Nonsynonymous SNP in ATG16L1 Predisposes to Ileal Crohn's Disease and Is Independent of CARD15 and IBD5. Gastroenterology 132, 1665–1671 (2007).

Raelson, J. V et al. Genome-wide association study for Crohn's disease in the Quebec Founder Population identifies multiple validated disease loci. Proceedings of the National Academy of Sciences of the United States of America 104, 14747–52 (2007).

Rosenstiel, P. et al. Functional characterization of two novel 5' untranslated exons reveals a complex regulation of NOD2 protein expression. BMC Genomics 8, 472 (2007).

Schafmayer, C. et al. Genetic investigation of DNA-repair pathway genes PMS2, MLH1, MSH2, MSH6, MUTYH, OGG1 and MTH1 in sporadic colon cancer. Int J Cancer 121, 555–558 (2007).

Schafmayer, C. et al. Investigation of the Lith6 candidate genes APOBEC1 and PPARG in human gallstone disease. Liver Int 27, 910–919 (2007).

Wollstein, A. et al. Efficacy assessment of SNP sets for genome-wide disease association studies. Nucleic Acids Res 35, e113 (2007).

Franke, A. et al. Replication of signals from recent studies of Crohn's disease identifies previously unknown disease loci for ulcerative colitis. *Nature genetics* 40, 713–5 (2008).

Franke, A. et al. Sequence variants in *IL10*, *ARPC2* and multiple other loci contribute to ulcerative colitis susceptibility. *Nature genetics* 40, 1319–23 (2008).

Franke, A. et al. Genome-wide association analysis in sarcoidosis and Crohn's disease unravels a common susceptibility locus on 10p12.2. *Gastroenterology* 135, 1207–1215 (2008).

Hofmann, S. et al. Genome-wide association study identifies *ANXA11* as a new susceptibility locus for sarcoidosis. *Nature genetics* 40, 1103–6 (2008).

Kaser, A. et al. *XBP1* links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease. *Cell* 134, 743–756 (2008).

Nothnagel, M., Ellinghaus, D., Schreiber, S., Krawczak, M. & Franke, A. A comprehensive evaluation of SNP genotype imputation. *Hum Genet* (2008).

Perminow, G. et al. Characterization in childhood inflammatory bowel disease. A new population-based inception cohort from South-Eastern Norway, 2005–07, showing increased incidence in Crohn's disease. *Scand J Gastroenterol* 44, 1–11 (2008).

Schafmayer, C. et al. Investigation of the colorectal cancer susceptibility region on chromosome 8q24.21 in a large German case-control sample. *Int J Cancer* 124, 75–80 (2008).

Dibbens, L. M. et al. Familial and sporadic 15q13.3 microdeletions in Idiopathic Generalized Epilepsy: Precedent for Disorders with Complex Inheritance. *Hum Mol Genet* (2009).doi:ddp311 [pii] 10.1093/hmg/ddp311

Ellinghaus, D., Schreiber, S., Franke, A. & Nothnagel, M. Current software for genotype imputation. *Hum Genomics* 3, 371–380 (2009).

End, C. et al. *DMBT1* functions as pattern-recognition molecule for poly-sulfated and poly-phosphorylated ligands. *Eur J Immunol* 39, 833–42 (2009).

Esparza-Gordillo, J. et al. A common variant on chromosome 11q13 is associated with atopic dermatitis. *Nature genetics* 41, 596–601 (2009).

Helbig, I. et al. 15Q13.3 Microdeletions Increase Risk of Idiopathic Generalized Epilepsy. *Nature genetics* 41, 160–2 (2009).

Henckaerts, L. et al. Genetic Risk Profiling And Prediction Of Disease Course In Crohn'S Disease Patients. *Clin Gastroenterol Hepatol* 7, 972–980.e2 (2009).

Lin, Z. et al. Genetic association of *DLG5* R30Q with familial and sporadic inflammatory bowel disease in men. *Dis Markers* 27, 193–201 (2009).

Loos, B. G. et al. *NOD1* gene polymorphisms in relation to aggressive periodontitis. *Innate Immun* 15, 225–32 (2009).

Melum, E., Franke, A. & Karlsen, T. H. Genome-wide association studies--a summary for the clinical gastroenterologist. *World J Gastroenterol* 15, 5377–5396 (2009).

Möckelmann, N. et al. Investigation of innate immunity genes *CARD4*, *CARD8* and *CARD15* as germline susceptibility factors for colorectal cancer. *BMC Gastroenterol* 9, 79 (2009).

Nair, R. P. et al. Genome-wide scan reveals association of psoriasis with *IL-23* and *NF-kappaB* pathways. *Nature genetics* 41, 199–204 (2009).

Nebel, A. et al. A functional *EXO1* promoter variant is associated with prolonged life expectancy in centenarians. *Mech Ageing Dev* 130, 691–699 (2009).

Rosenstiel, P., Sina, C., Franke, A. & Schreiber, S. Towards a molecular risk map--recent advances on the etiology of inflammatory bowel disease. *Seminars in immunology* 21, 334–45 (2009).

Sina, C. et al. G protein-coupled receptor 43 is essential for neutrophil recruitment during intestinal inflammation. *J Immunol* 183, 7514–7522 (2009).

Teuber, M., Wenz, M. H., Schreiber, S. & Franke, A. GMFilter and SXTTestPlate: software tools for improving the SNPlex genotyping system. *BMC Bioinformatics* 10, 81 (2009).

Weersma, R. K. et al. Confirmation of multiple Crohn's disease susceptibility loci in a large dutch-belgian cohort. *Am J Gastroenterol* 104, 630–638 (2009).

Yu, X. et al. Association of *UCP2* -866 G/A polymorphism with chronic inflammatory diseases. *Genes Immun* 10, 601–5 (2009).

Akil, I. et al. Co-existence of chronic renal failure, renal clear cell carcinoma, and Blau syndrome. *Pediatr Nephrol* 25, 977–981 (2010).

Buch, S. et al. Loci From a Genome-Wide Analysis of Bilirubin Levels Are Associated With Gallstone Risk and Composition. *Gastroenterology* (2010).doi:S0016-5085(10)01313-2 [pii] 10.1053/j.gastro.2010.09.003

Chapman, J. A. et al. The dynamic genome of *Hydra*. *Nature* 464, 592–596 (2010).

de Kovel, C. G. F. et al. Recurrent microdeletions at 15q11.2 and 16p13.11 predispose to idiopathic generalized epilepsies. *Brain* 133, 23–32 (2010).

Ding, J. et al. Gene expression in skin and lymphoblastoid cells: Refined statistical method reveals extensive overlap in cis-eQTL signals. *Am J Hum Genet* 87, 779–789 (2010).

Ellinghaus, E. et al. Genome-wide association study identifies a psoriasis susceptibility locus at *TRAF3IP2*. *Nature genetics* 42, 991–5 (2010).

Festen, E. A. M. et al. Genetic analysis in a Dutch study sample identifies more ulcerative colitis susceptibility loci and shows their additive role in disease risk. *The American journal of gastroenterology* 105, 395–402 (2010).

Flachsbart, F. et al. Investigation of genetic susceptibility factors for human longevity - a targeted nonsynonymous SNP study. *Mutat Res* 694, 13–19 (2010).

Franke, A. et al. Genome-wide association study for ulcerative colitis identifies risk loci at 7q22 and 22q13 (*IL17REL*). *Nature genetics* 42, 292–4 (2010).

Franke, A. et al. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nature genetics* 42, 1118–25 (2010).

Holm, K., Melum, E., Franke, A. & Karlsen, T. H. SNPexp - A web tool for calculating and visualizing correlation between HapMap genotypes and gene expression levels. *BMC Bioinformatics* 11, 600 (2010).

Horn, D. et al. Identification of FOXP1 deletions in three unrelated patients with mental retardation and significant speech and language deficits. *Hum Mutat* 31, E1851–60 (2010).

Hov, J. R. et al. Mutational characterization of the bile acid receptor TGR5 in primary sclerosing cholangitis. *PloS one* 5, e12403 (2010).

Juran, B. D. et al. Genetic polymorphisms of matrix metalloproteinase 3 in primary sclerosing cholangitis. *Liver Int* (2010).doi:10.1111/j.1478-3231.2010.02420.x

Karlsen, T. H. et al. Genome-wide association analysis in primary sclerosing cholangitis. *Gastroenterology* 138, 1102–11 (2010).

Karlsen, T. H., Melum, E. & Franke, A. The utility of genome-wide association studies in hepatology. *Hepatology (Baltimore, Md.)* 51, 1833–42 (2010).

Köttgen, A. et al. New loci associated with kidney function and chronic kidney disease. *Nature genetics* 42, 376–84 (2010).

Lin, Z. et al. OCTN1 variant L503F is associated with familial and sporadic inflammatory bowel disease. *J Crohns Colitis* 4, 132–138 (2010).

Lin, Z. et al. Genetic association of nonsynonymous variants of the IL23R with familial and sporadic inflammatory bowel disease in women. *Dig Dis Sci* 55, 739–746 (2010).

Liu, J. Z. et al. Meta-analysis and imputation refines the association of 15q25 with smoking quantity. *Nature genetics* 42, 436–40 (2010).

Lorentzen, A. R. et al. Association to the Glypican-5 gene in multiple sclerosis. *J Neuroimmunol* 226, 194–197 (2010).

Mefford, H. C. et al. Genome-wide copy number variation in epilepsy: novel susceptibility loci in idiopathic generalized and focal epilepsies. *PLoS Genet* 6, e1000962 (2010).

Melum, E. et al. SNP discovery performance of two second-generation sequencing platforms in the NOD2 gene region. *Hum Mutat* 31, 875–885 (2010).

Möller, M. et al. A functional haplotype in the 3'untranslated region of TNFRSF1B is associated with tuberculosis in two African populations. *Am J Respir Crit Care Med* 181, 388–393 (2010).

Muhle, H. et al. Role of GRM4 in idiopathic generalized epilepsies analysed by genetic association and sequence analysis. *Epilepsy Res* 89, 319–326 (2010).

Muhle, H. et al. A duplication in 1q21.3 in a family with early onset and childhood absence epilepsy. *Epilepsia* 51, 2453–2456 (2010).

Noor, A. et al. Disruption at the *PTCHD1* Locus on Xp22.11 in Autism spectrum disorder and intellectual disability. *Sci Transl Med* 2, 49ra68 (2010).

Pattaro, C. et al. A meta-analysis of genome-wide data from five European isolates reveals an association of *COL22A1*, *SYT1*, and *GABRR2* with serum creatinine level. *BMC Med Genet* 11, 41 (2010).

Stuart, P. E. et al. Genome-wide association analysis identifies three psoriasis susceptibility loci. *Nature genetics* 42, 1000–4 (2010).

Sun, L.-D. et al. Association analyses identify six new psoriasis susceptibility loci in the Chinese population. *Nature genetics* 42, 1005–9 (2010).

Sventoraityte, J. et al. *NOD2*, *IL23R* and *ATG16L1* polymorphisms in Lithuanian patients with inflammatory bowel disease. *World J Gastroenterol* 16, 359–364 (2010).

Wittig, M., Helbig, I., Schreiber, S. & Franke, A. CNVneta: a data mining tool for large case-control copy number variation datasets. *Bioinformatics* 26, 2208–2209 (2010).

Andersen, V. et al. Assessment of heterogeneity between European Populations: a Baltic and Danish replication case-control study of SNPs from a recent European ulcerative colitis genome wide association study. *BMC Med Genet* 12, 139 (2011).

Anderson, C. A. et al. Meta-analysis identifies 29 additional ulcerative colitis risk loci, increasing the number of confirmed associations to 47. *Nature genetics* 43, 246–52 (2011).

Angelova-Fischer, I. et al. Distinct barrier integrity phenotypes in filaggrin-related atopic eczema following sequential tape stripping and lipid profiling. *Exp Dermatol* 20, 351–356 (2011).

Del Greco, M. F. et al. Genome-wide association analysis and fine mapping of *NT-proBNP* level provide novel insight into the role of the *MTHFR-CLCN6-NPPA-NPPB* gene cluster. *Hum Mol Genet* 20, 1660–1671 (2011).

Dolmans, G. H. et al. Wnt signaling and Dupuytren's disease. *The New England journal of medicine* 365, 307–17 (2011).

Ehret, G. B. et al. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* 478, 103–109 (2011).

Ellinghaus, E. et al. Identification of germline susceptibility loci in *ETV6-RUNX1*-rearranged childhood acute lymphoblastic leukemia. *Leukemia* (2011).doi:leu2011302 [pii] 10.1038/leu.2011.302

Ellinghaus, E. et al. Genome-Wide Meta-Analysis of Psoriatic Arthritis Identifies Susceptibility Locus at *REL*. *J Invest Dermatol* (2011).doi:jid2011415 [pii] 10.1038/jid.2011.415

Eyerich, S. et al. Mutual antagonism of T cells causing psoriasis and atopic eczema. *N Engl J Med* 365, 231–238 (2011).

Fischer, A. et al. Association of inflammatory bowel disease risk loci with sarcoidosis, and its acute and chronic subphenotypes. *Eur Respir J* 37, 610–616 (2011).

Folseraas, T., Melum, E., Franke, A. & Karlsen, T. H. Genetics in primary sclerosing cholangitis. *Best practice & research. Clinical gastroenterology* 25, 713–26 (2011).

Freiwald, A. et al. Differential analysis of Crohn's disease and ulcerative colitis by mass spectrometry. *Inflamm Bowel Dis* 17, 1051–1052 (2011).

Gieger, C. et al. New gene functions in megakaryopoiesis and platelet formation. *Nature* 480, 201–208 (2011).

Hofmann, S. et al. A genome-wide association study reveals evidence of association with sarcoidosis at 6p12.1. *Eur Respir J* (2011).doi:09031936.00001711 [pii] 10.1183/09031936.00001711

Hunt, K. A. et al. Rare and functional SIAE variants are not associated with autoimmune disease risk in up to 66,924 individuals of European ancestry. *Nature genetics* 44, 3–5 (2011).

Jacobs, G. et al. Polymorphisms in the 3'UTR of the CDH1 gene are a risk factor for primary gastric diffuse large B-cell lymphoma. *Haematologica* (2011).doi:haematol.2010.033126 [pii] 10.3324/haematol.2010.033126

Janse, M. et al. Three ulcerative colitis susceptibility loci are associated with primary sclerosing cholangitis and indicate a role for IL2, REL, and CARD9. *Hepatology* 53, 1977–1985 (2011).

Keller, A. et al. Toward the blood-borne miRNome of human diseases. *Nat Methods* (2011).doi:nmeth.1682 [pii] 10.1038/nmeth.1682

Kirschner, J. et al. Mapping of quantitative trait loci controlling lifespan in the short-lived fish *Nothobranchius furzeri*- a new vertebrate model for age research. *Aging Cell* 49, no–no (2011).

Klostermeier, U. C. et al. A tissue-specific landscape of sense/antisense transcription in the mouse intestine. *BMC Genomics* 12, 305 (2011).

Kreck, B. et al. B-SOLANA: An approach for the analysis of two-base encoding bisulfite sequencing data. *Bioinformatics (Oxford, England)* 28, 428–429 (2011).

Lionel, A. C. et al. Rare copy number variation discovery and cross-disorder comparisons identify risk genes for ADHD. *Sci Transl Med* 3, 95ra75 (2011).

Melum, E. et al. Genome-wide association analysis in primary sclerosing cholangitis identifies two non-HLA susceptibility loci. *Nature genetics* 43, 17–9 (2011).

Mertes, F. et al. Targeted enrichment of genomic DNA regions for next-generation sequencing. *Briefings in functional genomics* 10, 374–86 (2011).

Naumann, A. et al. A comprehensive analysis of the COL29A1 gene does not support a role in eczema. *J Allergy Clin Immunol* 127, 1187–94 e7 (2011).

Nebel, A. et al. A genome-wide association study confirms APOE as the major gene influencing survival in long-lived individuals. *Mech Ageing Dev* 132, 324–330 (2011).

Paternoster, L. et al. Meta-analysis of genome-wide association studies identifies three new risk loci for atopic dermatitis. *Nature genetics* (2011).doi:10.1038/ng.1017

Rausch, P. et al. Colonic mucosa-associated microbiota is influenced by an interaction of Crohn disease and FUT2 (Secretor) genotype. *Proc Natl Acad Sci U S A* 108, 19030–19035 (2011).

Rieder, F. et al. Characterization of changes in serum anti-glycan antibodies in Crohn's disease - a longitudinal analysis. *PLoS ONE* 6, e18172 (2011).

Rivas, M. A. et al. Deep resequencing of GWAS loci identifies independent rare variants associated with inflammatory bowel disease. *Nature genetics* (2011).doi:10.1038/ng.952

Rossin, E. J. et al. Proteins encoded in genomic regions associated with immune-mediated disease physically interact and suggest underlying biology. *PLoS genetics* 7, e1001273 (2011).

Sanson, M. et al. Chromosome 7p11.2 (EGFR) variation influences glioma risk. *Hum Mol Genet* (2011).doi:ddr192 [pii] 10.1093/hmg/ddr192

Sawcer, S. et al. Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature* 476, 214–219 (2011).

Sun, L.-D. et al. Genome-wide association study identifies two new susceptibility loci for atopic dermatitis in the Chinese Han population. *Nature genetics* 43, 690–4 (2011).

von Spiczak, S. et al. Comprehensive analysis of candidate genes for photosensitivity using a complementary bioinformatic and experimental approach. *Epilepsia* 52, e143–7 (2011).

Wellge, B. E. et al. Pregnancy in primary sclerosing cholangitis. *Gut* (2011).doi:gut.2010.228924 [pii] 10.1136/gut.2010.228924

Dizier, M.-H. et al. The ANO3/MUC15 locus is associated with eczema in families ascertained through asthma. *The Journal of allergy and clinical immunology* 129, 1547–1553.e3 (2012).

Ellinghaus, D. et al. Combined analysis of genome-wide association studies for Crohn disease and psoriasis identifies seven shared susceptibility loci. *American journal of human genetics* 90, 636–47 (2012).

ElSharawy, A. et al. Improving mapping and SNP-calling performance in multiplexed targeted next-generation sequencing. *BMC Genomics* 13, 417 (2012).

Elsharawy, A. et al. Genome-wide miRNA signatures of human longevity. *Aging cell* 1–10 (2012).doi:10.1111/j.1474-9726.2012.00824.x

ElSharawy, A. et al. Accurate variant detection across non-amplified and whole genome amplified DNA using targeted next generation sequencing. *BMC Genomics* 13, 500 (2012).

Folseraas, T. et al. Extended Analysis of a Genome-Wide Association Study in Primary Sclerosing Cholangitis Detects Multiple Novel Risk Loci. *Journal of hepatology* xxx, 1–10 (2012).

Forster, M. et al. From next-generation sequencing alignments to accurate comparison and validation of single-nucleotide variants: the pibase software. *Nucleic Acids Research* 1–12 (2012).doi:10.1093/nar/gks836

Häsler, R. et al. A functional methylome map of ulcerative colitis. *Genome research* (2012).doi:10.1101/gr.138347.112

Helbig, K. L. et al. A case-only study of gene-environment interaction between genetic susceptibility variants in *NOD2* and cigarette smoking in Crohn's disease aetiology. *BMC medical genetics* 13, 14 (2012).

Huang, J., Ellinghaus, D., Franke, A., Howie, B. & Li, Y. 1000 Genomes-based imputation identifies novel and refined associations for the Wellcome Trust Case Control Consortium phase 1 Data. *European journal of human genetics* : *EJHG* 1–5 (2012).doi:10.1038/ejhg.2012.3

Invernizzi, P. et al. Classical *HLA-DRB1* and *DPB1* alleles account for HLA associations with primary biliary cirrhosis. *Genes and immunity* 1–8 (2012).doi:10.1038/gene.2012.17

Keller, A. et al. New insights into the Tyrolean Iceman's origin and phenotype as inferred by whole-genome sequencing. *Nature communications* 3, 698 (2012).

Krueger, F., Kreck, B., Franke, A. & Andrews, S. R. DNA methylome analysis using short bisulfite sequencing data. *Nature methods* 9, 145–51 (2012).

Næss, S., Shiryaev, A., Hov, J. R., Franke, A. & Karlsen, T. H. Genetics in primary sclerosing cholangitis. *Clinics and research in hepatology and gastroenterology* (2012).doi:10.1016/j.clinre.2012.02.003

Olszak, T. et al. Microbial Exposure During Early Life Has Persistent Effects on Natural Killer T Cell Function. *Science* (New York, N.Y.) 6–10 (2012).doi:10.1126/science.1219328

Pattaro, C. et al. Genome-wide association and functional follow-up reveals new loci for kidney function. *PLoS genetics* 8, e1002584 (2012).

Rebane, A. et al. Mechanisms of IFN- γ -induced apoptosis of human skin keratinocytes in patients with atopic dermatitis. *The Journal of allergy and clinical immunology* 1, 1297–1306 (2012).

Srivastava, B. et al. Fine mapping and replication of genetic risk loci in primary sclerosing cholangitis. *Scandinavian journal of gastroenterology* 820–826 (2012).doi:10.3109/00365521.2012.682090

Yu, W. et al. *PTPN2* is associated with Crohn's disease and its expression is regulated by *NKX2-3*. *Disease markers* 32, 83–91 (2012).